

# Generalized Electromagnetic fields in Chiral Medium

P. S. Bisht<sup>1</sup>, Jivan Singh<sup>2</sup> and O. P. S. Negi<sup>1</sup>

February 2, 2008

<sup>1</sup>Department of Physics  
Kumaun University  
Soban Singh Jeena Campus  
Almora - 263601 (Uttarakhand)  
INDIA

<sup>2</sup>Department of Physics  
Govt. Post Graduate College  
Pithoragarh (Uttarakhand)  
INDIA

Email:- ps\_bisht123@rediffmail.com  
jgaria@indiatimes.com  
ops\_negi@yahoo.co.in

## Abstract

The time dependent Dirac-Maxwell's Equations in presence of electric and magnetic sources are written in chiral media and the solutions for the classical problem are obtained in unique simple and consistent manner. The quaternion reformulation of generalized electromagnetic fields in chiral media has also been developed in compact, simple and consistent manner.

# 1 Introduction

Physicists were fascinated about magnetic monopoles since its ingenious idea was put forward by Dirac [1] and also by Saha [2]. So many attempts [3, 4] were made for the experimental verification of conclusive existence of magnetic monopoles and after the failure of attempts, the literature [5, 6, 7] turned partially negative casting doubts on the existence of such particles. The work of the Schwinger [8] was the first exception to the argument against the existence of monopoles. At the same time so many paradoxes were related to the theory of pure Abelian monopoles, as Dirac's veto, wrong spin-statistics connection [9] and many others [10, 11]. Several problems were soon resolved by the invention of dyons [12, 13, 14, 15] particles carrying simultaneous existence of electric and magnetic charges. Fresh interest in this subject was enhanced by the idea given by t'Hooft [16] and Polyakov [17] showing that monopoles are the intrinsic parts of grand unified theories. The Dirac monopole is an elementary particle but the t'Hooft-Polyakov monopole [16, 17] is a complicated extended object having a definite mass and finite size inside of which massive fields play a role in providing a smooth structure and outside it they vanish rapidly leaving the field configuration identical to Abelian Dirac monopole. Julia and Zee [178] have extended the idea of t' Hooft [16] and Polyakov [17] to construct the classical solutions for non-Abelian dyon. On the other hand, Prasad and Sommerfield [19, 20] have derived the analytic stable solutions for the non-Abelian monopoles and dyons of finite mass by keeping the symmetry of vacuum broken but letting the self-interaction of Higgs field approaching zero. Such solutions, satisfying Bogomolnys condition [21] are described as Bogomolny-Prasad-Sommerfield (BPS) monopoles. Kranchenko and coauthors [22, 23] discussed the Maxwell's equations in homogenous media and developed [24] the quaternionic reformulation of the time-dependent Maxwell's equations along with the classical solution of a moving source i.e. electron. They have also demonstrated [25] the electromagnetic fields in chiral media along with their quaternionic reformulation in a simple and consistent manner. Recently, we have extended the work of Kranchenko [22, 23] and reformulated [26] the Maxwell's -Dirac equation in homogenous (isotropic) medium and their quaternionic forms in a unique and consistent way [27]. We have also described [28] the time-harmonic Maxwell's equations for generalized fields of dyons in a consistent manner. Keeping these facts in mind, in this paper, we have derived the generalized theory of Maxwell's -Dirac equation in presence of electric and magnetic charges in chiral and homogenous

(isotropic) medium. The quaternion analysis of these time dependent Maxwell's equations in presence of electric and magnetic charges are obtained in unique, simpler and consistent manner. It has been shown that the quantum equations and equation of motion represent the dynamics of electric charge similar to the theory described by Kranchenko [22, 23] in the absence of magnetic monopole or vice versa.

## 2 Generalized Maxwell's Equation of Dyons in Isotropic Medium

Considering the existence of magnetic monopoles, Dirac [1] generalized the Maxwell's field equations in homogenous (isotropic) medium in the following manner [26];

$$\begin{aligned}
\vec{\nabla} \cdot \vec{E} &= \frac{\rho_e}{\epsilon} \\
\vec{\nabla} \cdot \vec{B} &= \mu \rho_m \\
\vec{\nabla} \times \vec{E} &= -\frac{\partial \vec{B}}{\partial t} - \frac{\vec{j}_m}{\epsilon} \\
\vec{\nabla} \times \vec{B} &= \frac{1}{v^2} \frac{\partial \vec{E}}{\partial t} + \mu \vec{j}_e
\end{aligned} \tag{1}$$

where  $\rho_e$  and  $\rho_m$  are respectively the electric and magnetic charge densities while  $\vec{j}_e$  and  $\vec{j}_m$  are the corresponding current densities,  $\vec{D}$  is electric induction vector,  $\vec{E}$  is electric field,  $\vec{B}$  is magnetic field and  $\epsilon_r$  and  $\mu_r$  are defined respectively as relative permittivity and permeability in electric and magnetic fields.

The electric and magnetic fields of dyons are expressed in homogenous medium in terms of two potentials as,

$$\vec{E} = -\vec{\nabla} \phi_e - \frac{\partial \vec{C}}{\partial t} - \vec{\nabla} \times \vec{D} \tag{2}$$

$$\vec{B} = -\vec{\nabla} \phi_m - \frac{1}{v^2} \frac{\partial \vec{D}}{\partial t} + \vec{\nabla} \times \vec{C} \tag{3}$$

where  $\{C^\mu\} = \{\phi_e, v \vec{C}\}$  and  $\{D^\mu\} = \{v \phi_m, \vec{D}\}$  are the two four-potentials associated with electric and magnetic charges. Let us define the complex vector

field  $\vec{\psi}$  in the following form

$$\vec{\psi} = \vec{E} - i v \vec{B}. \quad (4)$$

Equations (2,3) and (4) establish the following relation between generalized field vector and the components of generalized four-potential as,

$$\vec{\psi} = -\frac{\partial \vec{V}}{\partial t} - \vec{\nabla} \phi - i v (\vec{\nabla} \times \vec{V}) \quad (5)$$

where  $\{V_\mu\}$  is the generalised four-potential of dyons in homogenous medium and is defined as

$$V_\mu = \{\phi, \vec{V}\} \quad (6)$$

i.e.

$$\phi = \phi_e - i v \phi_m \quad (7)$$

and

$$\vec{V} = \vec{C} - i \frac{\vec{D}}{v}. \quad (8)$$

Maxwell's field equation (1) may then be written in terms of generalized field  $\vec{\psi}$  as

$$\vec{\nabla} \cdot \vec{\psi} = \frac{\rho}{\epsilon} \quad (9)$$

$$\vec{\nabla} \times \vec{\psi} = -i v (\mu \vec{J} + \frac{1}{v^2} \frac{\partial \vec{\psi}}{\partial t}) \quad (10)$$

where  $\rho$  and  $\vec{J}$  the generalized charge and current source densities of dyons in homogenous medium given by

$$\rho = \rho_e - i \frac{\rho_m}{v} \quad (11)$$

$$\vec{J} = \vec{j}_e - iv \vec{j}_m. \quad (12)$$

Using equation (9) we introduce a new parameter  $\vec{S}$  (i.e. the field current) as

$$\vec{S} = \square \vec{\psi} = -\mu \frac{\partial \vec{j}}{\partial t} - \frac{1}{\epsilon} \vec{\nabla} \rho - iv \mu (\vec{\nabla} \times \vec{j}) \quad (13)$$

where  $\square$  is the D'Alembertian operator.

In terms of components of complex potential the Maxwell's -Dirac equations are written as

$$\square \phi = v \mu \rho \quad (14)$$

$$\square \vec{V} = \mu \vec{J}. \quad (15)$$

We may thus write the tensorial form of generalized Maxwell's -Dirac equations of dyons in homogenous medium as

$$F_{\mu\nu,\nu} = j_\mu^e \quad (16)$$

$$F_{\mu\nu,\nu}^d = j_\mu^m. \quad (17)$$

Defining generalized field tensor of dyon as

$$G_{\mu\nu} = F_{\mu\nu} - iv F_{\mu\nu}^d \quad (18)$$

which yields the following covariant form of generalized field equation of dyon in homogenous (isotropic) medium i.e.

$$G_{\mu\nu,\nu} = J_\mu \quad (19)$$

$$G_{\mu\nu,\nu}^d = -i J_\mu^* . \quad (20)$$

### 3 Generalized Maxwell's Equation for Homogeneous Medium in Quaternionic Form

A quaternion is defined as

$$q = q_0 e_0 + q_1 e_1 + q_2 e_2 + q_3 e_3 \quad (21)$$

where  $q_0, q_1, q_2, q_3$  are real numbers and called the components of the quaternion  $q$  and the quaternion units  $e_0, e_1, e_2, e_3$  satisfy the following multiplication rules;

$$\begin{aligned} e_0^2 &= 1 \\ e_j e_k &= -\delta_{jk} + \epsilon_{jkl} e_l \end{aligned} \quad (22)$$

where  $\delta_{jk}$  and  $\epsilon_{jkl}$  ( $j, k, l = 1, 2, 3$  and  $e_0 = 1$ ) are respectively the Kronecker delta and three-index Levi-Civita symbol. For any quaternion, there exists a quaternion conjugate

$$\bar{q} = q - q_1 e_1 - q_2 e_2 - q_3 e_3 = q_0 - \vec{q}. \quad (23)$$

Quaternion conjugate is an automorphism of ring of quaternion i.e.

$$(\overline{pq}) = (\bar{q})(\bar{p}). \quad (24)$$

The norm of a quaternion is given as

$$N(q) = q \cdot \bar{q} = \bar{q} \cdot q = q_0^2 + q_1^2 + q_2^2 + q_3^2 = |q|^2. \quad (25)$$

The inverse of a quaternion is also a quaternion

$$q^{-1} = \frac{\bar{q}}{|q|^2}. \quad (26)$$

The quaternionic form of differential operator may be defined as [27]

$$\boxed{\cdot} = \left(-\frac{i}{v}\partial_t + D\right)$$

and its quaternion conjugate as

$$\overline{\boxed{\cdot}} = \left(-\frac{i}{v}\partial_t - D\right) \quad (27)$$

where  $D = \partial_1 e_1 + \partial_2 e_2 + \partial_3 e_3$ . Defining the complex vector wave function of generalized electromagnetic field as [26]

$$\vec{\psi} = \vec{E} - iv\vec{B}. \quad (28)$$

We can now express the generalized four-potential, four current, electric field and magnetic in quaternionic form as [27] ;

$$V = -i\frac{\phi}{v} + V_1 e_1 + V_2 e_2 + V_3 e_3 \quad (29)$$

$$J = -i\rho v + J_1 e_1 + J_2 e_2 + J_3 e_3 \quad (30)$$

$$E = E_1 e_1 + E_2 e_2 + E_3 e_3 \quad (31)$$

$$B = B_1 e_1 + B_2 e_2 + B_3 e_3 \quad (32)$$

Operating first equation of (27) to equation (28) and using equation (22 ), we get

$$\overline{\boxed{\cdot}}\psi = J. \quad (33)$$

Similarly we operate first equation of (27) to equation (30) and using equation (22) , we get

$$\square J = S. \quad (34)$$

Equations (33) and (34) are the quaternion forms of field equations associated respectively with the generalized potential and currents of dyons in homogeneous isotropic medium. These equations are invariant under quaternion transformations as well as with homogeneous Lorentz transformations. Instead of four sets of differential equations of field equations we may write only one set of quaternion differential equations in compact ,simple and consistent manner. We may reinterpret our results as that the algebra over the field of real numbers corresponds four sets of differential equations given by equation (1) , the algebra over the field of complex numbers corresponds to two sets of differential equations given by equation (9-10) while the algebra over the field of quaternion variables corresponds only to one set of differential equations given by equation (33) . As such equations (33) and (34) are the quaternion forms of differential equations (9-10 and 13) and may be visualized as the quaternion reformulation of generalized potential and current of dyons in isotropic homogeneous medium in compact, simpler and consistent manner.

## 4 Generalized Electromagnetic Fields in Chiral Media

Let us consider the generalized Maxwell's equations for dyons in homogenous (isotropic) medium as

$$\begin{aligned} \vec{\nabla} \cdot \tilde{E}(x) &= \frac{\rho_e(x)}{\epsilon} \\ \vec{\nabla} \cdot \tilde{H}(x) &= \rho_m(x) \\ \vec{\nabla} \times \tilde{E}(x) &= -\frac{\tilde{j}_m(x)}{\epsilon} + i\omega \tilde{B}(x) \\ \vec{\nabla} \times \tilde{H}(x) &= \mu \tilde{j}_e(x) - i\omega \tilde{D}(x). \end{aligned} \quad (35)$$

In the above equation we have taken the electric field  $\vec{E}$  and magnetic field  $\vec{B}$  is time harmonic. Considering electric and magnetic field as under [28]

$$\begin{aligned}\vec{E}(x, t) &= \text{Re}(\vec{E}(x)e^{-i\omega t}) \\ \vec{B}(x, t) &= \text{Re}(\vec{B}(x)e^{-i\omega t}).\end{aligned}\quad (36)$$

Here we consider the Born-Fedorov constitutive equations [29, 30, 31, 32],

$$\begin{aligned}\tilde{D}(x) &= \epsilon(\tilde{E}(x) + \beta \nabla \times \tilde{E}(x)) \\ \tilde{B}(x) &= \mu(\tilde{H}(x) + \beta \nabla \times \tilde{H}(x))\end{aligned}\quad (37)$$

where  $\epsilon, \mu$ , and  $\beta$  are permittivity, permeability and chiral parameter respectively. If the medium is isotropic, the cartesian field components are given by

$$\begin{aligned}\tilde{D}_x &= \epsilon \tilde{E}_x + \epsilon \beta \left( \frac{\partial \tilde{E}_z}{\partial y} - \frac{\partial \tilde{E}_y}{\partial z} \right) \\ \tilde{D}_y &= \epsilon \tilde{E}_y + \epsilon \beta \left( \frac{\partial \tilde{E}_x}{\partial z} - \frac{\partial \tilde{E}_z}{\partial x} \right) \\ \tilde{D}_z &= \epsilon \tilde{E}_z + \epsilon \beta \left( \frac{\partial \tilde{E}_y}{\partial x} - \frac{\partial \tilde{E}_x}{\partial y} \right) \\ \tilde{B}_x &= \mu \tilde{B}_x + \mu \beta \left( \frac{\partial \tilde{B}_z}{\partial y} - \frac{\partial \tilde{B}_y}{\partial z} \right) \\ \tilde{B}_y &= \mu \tilde{B}_y + \mu \beta \left( \frac{\partial \tilde{B}_x}{\partial z} - \frac{\partial \tilde{B}_z}{\partial x} \right) \\ \tilde{B}_z &= \mu \tilde{B}_z + \mu \beta \left( \frac{\partial \tilde{B}_y}{\partial x} - \frac{\partial \tilde{B}_x}{\partial y} \right).\end{aligned}\quad (38)$$

Applying the condition given by (37) into third and fourth equation of equation (36) we get,

$$\begin{aligned}\nabla \times \tilde{E}(x) &= -\frac{\tilde{j}_m(x)}{\epsilon} + i\omega\mu(\tilde{H}(x) + \beta \nabla \times \tilde{H}(x)) \\ \nabla \times \tilde{H}(x) &= -\tilde{j}_e(x) - i\omega\epsilon(\tilde{E}(x) + \beta \nabla \times \tilde{E}(x)).\end{aligned}\quad (39)$$

If we introduce the following notations

$$\begin{aligned}
\widetilde{E}(x) &= -\sqrt{\mu}\vec{E}(x) \\
\widetilde{H}(x) &= \sqrt{\varepsilon}\vec{H}(x) \\
\widetilde{j_e}(x) &= \sqrt{\varepsilon}\vec{j_e}(x) \\
\widetilde{j_m}(x) &= \sqrt{\varepsilon}\vec{j_m}(x),
\end{aligned} \tag{40}$$

the first and second equation of equation (36) and equation (39) reduce to the following differential equations as,

$$\begin{aligned}
\nabla \cdot \vec{E} &= -\frac{\rho_e(x)}{\varepsilon\sqrt{\mu}} \\
\nabla \cdot \vec{H} &= \frac{\rho_m(x)}{\sqrt{\varepsilon}}
\end{aligned}$$

$$\begin{aligned}
\nabla \times \vec{E}(x) &= -\frac{\vec{j_m}}{\sqrt{\mu\varepsilon}} - i\alpha(\vec{H}(x) + \beta\nabla \times \vec{H}(x)) \\
\nabla \times \vec{H}(x) &= \vec{j_e}(x) + i\alpha(\vec{E}(x) + \beta\nabla \times \vec{E}(x))
\end{aligned} \tag{41}$$

where  $\alpha = \frac{\omega}{v}$  is denoted as the wave number.

## 5 Maxwell's Equation for Dyons in Quaternionic Chiral Media

Let us consider the following purely vectorial biquaternionic functions as,

$$\vec{m}(x) = \vec{E}(x) + i\vec{H}(x) \tag{42}$$

$$\vec{n}(x) = \vec{E}(x) - i\vec{H}(x). \tag{43}$$

Operating the quaternionic differential operator  $D$  on equation (42), we get

$$D\vec{m}(x) = \frac{\rho(x)}{\varepsilon\sqrt{\mu}}(1 + \alpha\beta) - \alpha\vec{m}(x) - \alpha\beta D\vec{m}(x) + i\vec{j}(x). \tag{44}$$

Thus the complex quaternionic function  $\vec{m}(x)$  satisfies the following equation

$$(D + \frac{\alpha}{(1 + \alpha\beta)})\vec{m}(x) = \frac{\rho(x)}{\epsilon\sqrt{\mu}} + i\frac{\vec{j}(x)}{(1 + \alpha\beta)}. \quad (45)$$

By analogy of equation (45), we obtain the equation for  $\vec{n}$  as,

$$(D - \frac{\alpha}{(1 - \alpha\beta)})\vec{n}(x) = \frac{\rho(x)}{\epsilon\sqrt{\mu}} - i\frac{\vec{j}(x)}{(1 - \alpha\beta)}. \quad (46)$$

On using equation (35) and equation (41), the continuity equation is described in the following manner,

$$\begin{aligned} \text{div}\vec{j}_e &= i\frac{\alpha\rho_e}{\epsilon\sqrt{\mu}} \\ \text{div}\vec{j}_m &= i\frac{\alpha\rho_m}{v\sqrt{\epsilon}}. \end{aligned} \quad (47)$$

Introducing the notations

$$\begin{aligned} \alpha_1 &= \frac{\alpha}{(1 + \alpha\beta)} \\ \alpha_2 &= \frac{\alpha}{(1 - \alpha\beta)} \end{aligned} \quad (48)$$

and using equations (47) and (48), we may write the differential equations (45) and (46) in the following forms i.e.,

$$\begin{aligned} (D + \alpha_1)\vec{m}(x) &= -i\frac{\text{div}\vec{j}(x)}{\alpha} - i\frac{\vec{j}(x)\alpha_1}{\alpha} \\ &= -\frac{i}{\alpha}(\alpha_1\vec{j}(x) + \text{div}\vec{j}(x)) \\ (D - \alpha_2)\vec{n}(x) &= \frac{i}{\alpha}(\alpha_1\vec{j}(x) - \text{div}\vec{j}(x)). \end{aligned} \quad (49)$$

In the above equations if we put  $\beta = 0$ , we get the quaternionic form of the generalized Maxwell's equation of dyons in the absence of chiral medium. In general the wave numbers  $\alpha_1$  and  $\alpha_2$  are different and physically characterize the propagation of waves of opposing circular polarizations.

## 6 Discussion

In the foregoing analysis, the equations (37) showing the dependence of the electric displacement vector and the magnetic induction vector on the electric and magnetic fields, do not take into account the chirality of the medium. Instead, they depend upon only on the conductivity, electric permittivity and magnetic permeability of the medium. Chirality is described as the asymmetry in the molecular structure where a molecule is chiral if it cannot be superimposed onto its mirror image. Presence of chirality results in the rotation of electromagnetic fields and its observable, particular in the microwave range even for the case of a particle consisting electric and magnetic charges (i.e. a dyon). Such experimental observations may be used in physical chemistry to characterize molecular structure. The present theory of generalized electrodynamics of dyons leads the connection between the mechanical parameters with the chirality and dielectric properties of the brain tissue considered as a bioplasma . Hence the proposal for dyonic bioplasma is being considered. Our theory reduces to the theories described earlier [22, 23, 24, 25, 29, 30, 31, 32] for the case of electric charge in the absence of magnetic monopole on dyon and consequently the theories of pure monopole be described from duality in the absence of electric charge on dyons.

## References

- [1] Dirac, P. A .M, Proc. R. Soc. London Sec , **A 133** (1931) 60.
- [2] Saha M.N., Ind. J. Phys., **10** (1936) 145; Phys. Rev., **75** (1949) 1968.
- [3] Malkus M.V.R., Phys. Rev., **83** (1951) 899.
- [4] Goto E., H. M. Kolm and K. W. Ford, Phys. Rev. **132** (1963) 387.
- [5] Rosenbaum D., Phys. Rev., **147** (1966) 891.
- [6] Zwanziger D.; Phys. Rev. , **B137** (1965) 647.
- [7] Goldhaber A. S., Phys. Rev., **B140** (1965) 1407.
- [8] Schwinger J., Phys. Rev., **144** (1966) 1087;**151** (1966) 1055, 1048; Science, **165** (1969) 757.
- [9] Goldhaber A. S., Phys. Rev. Lett., **36** (1976) 1122.

- [10] Zwanziger D., Phys. Rev., **176** (1968) 1489.
- [11] Seo K., Phys. Lett., **B126** (1983) 201.
- [12] Faddeev L. D., Les Houches lecture, session 20, North Holland, (1976).
- [13] Coleman S., Lectures delivered at the Int. school of Subnuclear Phys, (1975).
- [14] Zwanziger D., Phys. Rev., **176** (1968) 1480.
- [15] Cabibbo N. and Ferrari E., Nuovo Cim., **23** (1962) 1147.
- [16] 't Hooft G. , Nucl. Phys., **B79** (1974) 276; **B138** (1978) 1.
- [17] Polyakov A. M., JETP Lett, **20** (1974) 194; Zh ETF Pis'ma **B79** (1974) 276; Phys. Lett. **B59** (1975) 82.
- [178] Julia B and Zee A, Phys Rev , **D21** (1980) 2940.
- [19] Prasad M., Comm. Math. Phys., **80** (1980) 137.
- [20] Prasad M and Sommerfield C, Phys Rev Lett. , **35** (1975) 760.
- [21] Bogomol'nyi E P, Sov. J. Nucl Phys, **24** (1976) 449.
- [22] Kravchenko, V.V., Applied Quaternionic Analysis, Research and Exposition in Mathematics, **28** (2003), Heldermann Verlage, Germany.
- [23] Kravchenko, V.V., Zeitschrift Fur Analysis und Anwenduhgen, **21** (2002) 21.
- [24] K. V. Khmelnytskaya and Kravchenko, V.V., Zeitschrift Fur Analysis und Anwenduhgen, **56** (2001), 53.
- [25] S. M. Grudsky, K. V. Khmelnytskaya and V. V. Kravchenko, J. Phys. A: Math. Gen., **37** (2004), 4641.
- [26] Jivan Singh, P. S. Bisht and O. P. S. Negi, "Generalized fields of dyons in Isotropic Medium", arXiv:hep-th/0611208 .
- [27] Jivan Singh, P. S. Bisht and O. P. S. Negi, "Quaternion Analysis of generalized fields of dyons in Isotropic Medium", arXiv:hep-th/0703083.
- [28] P. S. Bisht, Jivan Singh and O. P. S. Negi, "Quaternion Analyticity of Time-Harmonic Dyon Field Equations", arXiv:hep-th/0703107.

- [29] C. Athanasiadis, G. Roach and I. Stratis, Math. Nachr., **250** (2003), 3.
- [30] A. Lakhtakia, Beltrami Fields in Chiral Media, World Scientific, Singapore (1994).
- [31] I. V. Lindell, A. H. Sihvola, S. A. tretyakov and A. J. Viitanen, Electromagnetic waves in chiral and bi-isotropic media, Artech House Publisher, Boston, MA (1994).
- [32] T. S. Hector and Z. L. Mario, Facultad De Ingeniera, U.T.A. (Chile), **11** (2003), 3.